

BOGOMOLOV CONJECTURE FOR CURVES OF GENUS 2 OVER FUNCTION FIELDS

ATSUSHI MORIWAKI

ABSTRACT. In this note, we will show that Bogomolov conjecture holds for a non-isotrivial curve of genus 2 over a function field.

1. INTRODUCTION

Let k be an algebraically closed field, X a smooth projective surface over k , Y a smooth projective curve over k , and $f : X \rightarrow Y$ a generically smooth semistable curve of genus $g \geq 2$ over Y . Let K be the function field of Y , $\overline{K}$ the algebraic closure of K , and C the generic fiber of f . Let $j : C(\overline{K}) \rightarrow \text{Jac}(C)(\overline{K})$ be a morphism given by $j(x) = (2g - 2)x - \omega_C$ and $\|\cdot\|_{NT}$ the semi-norm arising from the Neron-Tate height pairing on $\text{Jac}(C)(\overline{K})$. We set

$$B_C(P; r) = \{x \in C(\overline{K}) \mid \|j(x) - P\|_{NT} \leq r\}$$

for $P \in \text{Jac}(C)(\overline{K})$ and $r \geq 0$, and

$$r_C(P) = \begin{cases} -\infty & \text{if } \#(B_C(P; 0)) = \infty, \\ \sup \{r \geq 0 \mid \#(B_C(P; r)) < \infty\} & \text{otherwise.} \end{cases}$$

Bogomolov conjecture claims that, if f is non-isotrivial, then $r_C(P)$ is positive for all $P \in \text{Jac}(C)(\overline{K})$. Even to say that $r_C(P) \geq 0$ for all $P \in \text{Jac}(C)(\overline{K})$ is non-trivial because it contains Manin-Mumford conjecture, which was proved by Raynaud. Further, it is well known that the above conjecture is equivalent to say the following.

Conjecture 1.1 (Bogomolov conjecture). If f is non-isotrivial, then

$$\inf_{P \in \text{Jac}(C)(\overline{K})} r_C(P) > 0.$$

Moreover, we can think the following effective version of Conjecture 1.1.

Conjecture 1.2 (Effective Bogomolov conjecture). In Conjecture 1.1, there is an effectively calculated positive number r_0 with

$$\inf_{P \in \text{Jac}(C)(\overline{K})} r_C(P) \geq r_0.$$

Date: October, 1995 (Second version).

We would like to express our thanks to Prof. Liu for his nice remarks.

In [2], we proved that, if f is non-isotrivial and the stable model of $f : X \rightarrow Y$ has only irreducible fibers, then Conjecture 1.2 holds. More precisely,

$$\inf_{P \in \text{Jac}(C)(\overline{K})} r_C(P) \geq \begin{cases} \sqrt{12(g-1)} & \text{if } f \text{ is smooth,} \\ \sqrt{\frac{(g-1)^3}{3g(2g+1)}} \delta & \text{otherwise,} \end{cases}$$

where δ is the number of singularities in singular fibers of $f : X \rightarrow Y$. In this note, we would like to show the following result.

Theorem 1.3. *If f is non-isotrivial and $g = 2$, then f is not smooth and*

$$\inf_{P \in \text{Jac}(C)(\overline{K})} r_C(P) \geq \sqrt{\frac{2}{135}} \delta.$$

2. NOTATIONS AND IDEAS

In this section, we use the same notations as in §1. Let $\omega_{X/Y}^a$ be the dualizing sheaf in the sense of admissible pairing. (For details concerning admissible pairing, see [5] or [2].) First we note the following theorem. (cf. [5, Theorem 5.6] or [2, Corollary 2.3])

Theorem 2.1. *If $(\omega_{X/Y}^a \cdot \omega_{X/Y}^a)_a > 0$, then*

$$\inf_{P \in \text{Jac}(C)(\overline{K})} r_C(P) \geq \sqrt{(g-1)(\omega_{X/Y}^a \cdot \omega_{X/Y}^a)_a},$$

where $(\cdot)_a$ is the admissible pairing.

From now on, we assume $g = 2$. By the above theorem, in order to get Theorem 1.3, we need to estimate $(\omega_{X/Y}^a \cdot \omega_{X/Y}^a)_a$. First of all, we can set

$$(2.2) \quad (\omega_{X/Y}^a \cdot \omega_{X/Y}^a)_a = (\omega_{X/Y} \cdot \omega_{X/Y}) - \sum_{y \in Y} e_y,$$

where e_y is the number coming from the Green function of $f^{-1}(y)$. This number depends on the configuration of $f^{-1}(y)$. So, let us consider the classification of semistable curves of genus 2. Let E be a semistable curve of genus 2 over k and E' the stable model of E , that is, E' is a curve obtained by contracting all (-2) -curves in E . It is well known that there are 7-types of stable curves of genus 2. Thus, we have the classification of semistable curves of genus 2 according to type of E' as in Table 1. (In Table 1, the symbol A_n for a node means that the dual graph of (-2) -curves over the node is same as A_n type graph.) The exact value of e_y can be found in Table 2 and will be calculated in §3.

Next we need to think an estimation of $(\omega_{X/Y} \cdot \omega_{X/Y})$ in terms of type of $f^{-1}(y)$. According to Ueno [4], there is the canonical section s of

$$H^0(Y, \det(f_*(\omega_{X/Y}))^{10})$$

such that $d_y = \text{ord}_y(s)$ for $y \in Y$ can be exactly calculated under the assumption that $\text{char}(k) \neq 2, 3, 5$. The result can be found in Table 2. Prof. Liu points out that by works of T. Saito [3] and Q. Liu [1], the value d_y in Table 2 still holds even if $\text{char}(k) \leq 5$.

Let δ_y be the number of singularities in $f^{-1}(y)$. Then, by Noether formula,

$$\deg(\det(f_*(\omega_{X/Y}))) = \frac{(\omega_{X/Y} \cdot \omega_{X/Y}) + \sum_{y \in Y} \delta_y}{12}.$$

On the other hand, by the definition of d_y ,

$$\sum_{y \in Y} d_y = 10 \deg(\det(f_*(\omega_{X/Y}))).$$

Thus, we have

$$(2.3) \quad (\omega_{X/Y} \cdot \omega_{X/Y}) = \sum_{y \in Y} \left(\frac{6}{5} d_y - \delta_y \right).$$

Hence, by (2.2) and (2.3),

$$(2.4) \quad (\omega_{X/Y}^a \cdot \omega_{X/Y}^a)_a = \sum_{y \in Y} \left(\frac{6}{5} d_y - \delta_y - e_y \right).$$

Therefore, using (2.4) and Table 2, we have the following theorem. (Note that an inequality

$$\frac{abc}{ab + bc + ca} \leq \frac{a + b + c}{9}$$

holds for all positive numbers a, b, c .)

Theorem 2.5. *If f is non-isotrivial, then f is not smooth and*

$$(\omega_{X/Y}^a \cdot \omega_{X/Y}^a)_a \geq \frac{2}{135} \delta,$$

where $\delta = \sum_{y \in Y} \delta_y$.

Note that non-smoothness of f can be easily derived from the fact that the moduli space M_2 of curves of genus 2 is an affine variety.

3. CALCULATION OF e_y

Let us start calculations of e_y . If the stable model of a fiber is irreducible, e_y is calculated in [2]. Thus it is sufficient to calculate e_y for II(a), IV(a,b), VI(a,b,c) and VII(a,b,c). In these cases, the stable model has two irreducible components. Let $f^{-1}(y) = C_1 + \cdots + C_n$ be the irreducible decomposition of $f^{-1}(y)$. We set

$$D_y = \sum_{i=1}^n (\omega_{X/Y} \cdot C_i) v_i,$$

where v_i is the vertex in G_y corresponding to C_i . Especially, we denote by P and Q corresponding vertexes to stable components. Then, $D_y = P + Q$. Let μ and g_μ be the measure and the Green function defined by D_y . In the same way as in the Proof of Theorem 5.1 in [2],

$$e_y = -g_\mu(D_y, D_y) + 4c(G_y, D_y),$$

where $c(G_y, D_y)$ is the constant coming from g_μ . By the definition of $c(G_y, D_y)$,

$$c(G_y, D_y) = g_\mu(P, P) + g_\mu(P, D_y).$$

Therefore, we have

$$e_y = 7g_\mu(P, P) - g_\mu(Q, Q) + 2g_\mu(P, Q).$$

Here claim:

Lemma 3.1. $g_\mu(P, P) = g_\mu(Q, Q)$. In particular,

$$e_y = 6g_\mu(P, P) + 2g_\mu(P, Q).$$


Proof. By the definition of $c(G_y, D_y)$,

$$c(G_y, D_y) = g_\mu(P, P) + g_\mu(P, P + Q) = g_\mu(Q, Q) + g_\mu(Q, P + Q).$$

Thus, we can see $g_\mu(P, P) = g_\mu(Q, Q)$. □

In the following, we will calculate e_y for each type II(a), IV(a,b), VI(a,b,c) and VII(a,b,c). First we present the dual graph of each type and then show its calculation.

Type II(a).



In this case, $\mu = \frac{\delta_P}{2} + \frac{\delta_Q}{2}$ by [5, Lemma 3.7]. We fix a coordinate $s : G \rightarrow [0, a]$ with $s(P) = 0$ and $s(Q) = a$. If we set

$$g(x) = -\frac{s(x)}{2} + \frac{a}{4},$$


then, $\Delta(g) = \delta_P - \mu$ and $\int_G g\mu = 0$. Thus, $g(x) = g_\mu(P, x)$. Hence

$$g_\mu(P, P) = \frac{a}{4} \quad \text{and} \quad g_\mu(P, Q) = -\frac{a}{4}.$$

Thus

$$e_y = 6g_\mu(P, P) + 2g_\mu(P, Q) = a.$$

Type IV(a,b).



We fix coordinates $s : G_1 \rightarrow [0, a]$ and $t : G_2 \rightarrow [0, b]$ with $s(P) = 0$, $s(Q) = a$ and $t(Q) = 0$. In this case, $\mu = \frac{\delta_P}{2} + \frac{dt}{2b}$ by [5, Lemma 3.7]. We set

$$g(x) = \begin{cases} -\frac{s(x)}{2} + \frac{b+12a}{48} & \text{if } x \in G_1, \\ \frac{1}{2} \left(\frac{t(x)^2}{2b} - \frac{t(x)}{2} \right) + \frac{b-12a}{48} & \text{if } x \in G_2. \end{cases}$$


Then, g is continuous, $\Delta(g|_{G_1}) = \frac{\delta_P}{2} - \frac{\delta_Q}{2}$, and $\Delta(g|_{G_2}) = \frac{\delta_Q}{2} - \frac{dt}{2b}$. Thus, $\Delta(g) = \delta_P - \mu$. Moreover, $\int_G g\mu = 0$. Therefore, $g(x) = g_\mu(P, x)$. Hence

$$g_\mu(P, P) = \frac{b+12a}{48} \quad \text{and} \quad g_\mu(P, Q) = \frac{b-12a}{48}.$$

Thus

$$e_y = 6g_\mu(P, P) + 2g_\mu(P, Q) = a + \frac{b}{6}.$$

Type VI(a,b,c).



We fix coordinates $s : G_1 \rightarrow [0, a]$, $t : G_2 \rightarrow [0, b]$ and $u : G_3 \rightarrow [0, c]$ with $s(P) = 0$, $s(Q) = a$, $t(P) = 0$ and $u(Q) = 0$. In this case, $\mu = \frac{dt}{2b} + \frac{du}{2c}$ by [5, Lemma 3.7]. We set

$$g(x) = \begin{cases} \frac{1}{2} \left(\frac{t(x)^2}{2b} - \frac{t(x)}{2} \right) + \frac{b+c+12a}{48} & \text{if } x \in G_2, \\ -\frac{s(x)}{2} + \frac{b+c+12a}{48} & \text{if } x \in G_1, \\ \frac{1}{2} \left(\frac{u(x)^2}{2c} - \frac{u(x)}{2} \right) + \frac{b+c-12a}{48} & \text{if } x \in G_3. \end{cases}$$


Then, g is continuous, $\Delta(g|_{G_1}) = \frac{\delta_P}{2} - \frac{\delta_Q}{2}$, $\Delta(g|_{G_2}) = \frac{\delta_P}{2} - \frac{dt}{2b}$, and $\Delta(g|_{G_3}) = \frac{\delta_Q}{2} - \frac{du}{2c}$. Thus, $\Delta(g) = \delta_P - \mu$. Moreover, $\int_G g\mu = 0$. Therefore, $g(x) = g_\mu(P, x)$. Hence

$$g_\mu(P, P) = \frac{b+c+12a}{48} \quad \text{and} \quad g_\mu(P, Q) = \frac{b+c-12a}{48}.$$

Thus

$$e_y = 6g_\mu(P, P) + 2g_\mu(P, Q) = a + \frac{b+c}{6}.$$

Type VII(a,b,c).



We fix coordinates $s : G_1 \rightarrow [0, a]$, $t : G_2 \rightarrow [0, b]$ and $u : G_3 \rightarrow [0, c]$ with $s(P) = 0$, $s(Q) = a$, $t(P) = 0$, $t(Q) = b$, $u(P) = 0$ and $u(Q) = c$. In this case, $\mu = \frac{ds}{3a} + \frac{dt}{3b} + \frac{du}{3c}$ by [5, Lemma 3.7]. We set

$$g(x) = \begin{cases} \frac{s(x)^2}{6a} - \left(\frac{1}{6} + \frac{1}{2} \frac{bc}{ab+bc+ca} \right) s(x) + \frac{a+b+c}{108} + \frac{1}{4} \frac{abc}{ab+bc+ca} & \text{if } x \in G_1, \\ \frac{t(x)^2}{6b} - \left(\frac{1}{6} + \frac{1}{2} \frac{ac}{ab+bc+ca} \right) t(x) + \frac{a+b+c}{108} + \frac{1}{4} \frac{abc}{ab+bc+ca} & \text{if } x \in G_2, \\ \frac{u(x)^2}{6c} - \left(\frac{1}{6} + \frac{1}{2} \frac{ab}{ab+bc+ca} \right) u(x) + \frac{a+b+c}{108} + \frac{1}{4} \frac{abc}{ab+bc+ca} & \text{if } x \in G_3. \end{cases}$$

Then, g is continuous and $\Delta(g) = \Delta(g|_{G_1}) + \Delta(g|_{G_2}) + \Delta(g|_{G_3}) = \delta_P - \mu$. Moreover, $\int_G g\mu = 0$. Therefore, $g(x) = g_\mu(P, x)$. Hence






$$e_y = 6g_\mu(P, P) + 2g_\mu(P, Q) = \frac{2}{27}(a+b+c) + \frac{abc}{ab+bc+ca}.$$

REFERENCES

- [1] Q. Liu, Conducteur et discriminant minimal de courbes de genre 2, *Comp. Math.*, 94 (1994), 51–79.
- [2] A. Moriwaki, Bogomolov conjecture over function fields for stable curves with only irreducible fibers, preprint.
- [3] T. Saito, The discriminants of curves of genus 2, *Comp. Math.*, 69 (1989), 229–240.
- [4] K. Ueno, Discriminants of curves of genus 2 and arithmetic surfaces, *Algebraic Geometry and Commutative Algebra in Honor of Masaki NAGATA*, (1987), 749–770.
- [5] S. Zhang, Admissible pairing on a curve, *Invent. Math.*, 112 (1993), 171–193.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, KYOTO UNIVERSITY, KYOTO, 606-01, JAPAN
E-mail address: moriwaki@kum.kyoto-u.ac.jp

TABLE 1. Classification of semistable curve E of genus 2

Type of E	Description of the stable model E' of E	Figure of E' and types of singularities by contracting (-2) -curves in E
I	a smooth curve of genus 2	 $g = 2$
II(a)	two elliptic curves meeting at one point transversally	 $g = 1$ $g = 1$
III(a)	an elliptic curve with one node	 $g = 1$
IV(a,b)	a smooth elliptic curve and a rational curve with one node, which meet at one point transversally	 $g = 0$ $g = 1$
V(a,b)	a rational curve with two nodes	 $g = 0$



VI(a,b,c)	two rational curves with one node, which meet at one point transversally	 $g = 0$ $g = 0$
VII(a,b,c)	two smooth rational curves, which meet at three points transversally	 $g = 0$ $g = 0$

TABLE 2. δ_y , d_y and e_y

Type	δ_y	d_y	e_y
I	0	0	0
II(a)	a	$2a$	a
III(a)	a	a	$\frac{a}{6}$
IV(a,b)	$a + b$	$2a + b$	$a + \frac{b}{6}$
V(a, b)	$a + b$	$a + b$	$\frac{a + b}{6}$
VI(a,b,c)	$a + b + c$	$2a + b + c$	$a + \frac{b + c}{6}$
VII(a,b,c)	$a + b + c$	$a + b + c$	$\frac{2}{27}(a + b + c) + \frac{abc}{ab + bc + ca}$